

A Feature-based Constraint System for Logic Programming with Entailment

Hassan Aït-Kaci* Andreas Podelski* Gert Smolka†

Abstract

This paper presents the constraint system FT , which we feel is an intriguing alternative to Herbrand both theoretically and practically. As does Herbrand, FT provides a universal data structure based on trees. However, the trees of FT (called feature trees) are more general than the trees of Herbrand (called constructor trees), and the constraints of FT are finer grained and of different expressivity. The basic notion of FT are functional attributes called features, which provide for record-like descriptions of data avoiding the overspecification intrinsic in Herbrand’s constructor-based descriptions. The feature tree structure fixes an algebraic semantics for FT . We will also establish a logical semantics, which is given by three axiom schemes fixing the first-order theory FT .

FT is a constraint system for logic programming, providing a test for unsatisfiability, and a test for entailment between constraints, which is needed for advanced control mechanisms.

The two major technical contributions of this paper are (1) an incremental entailment simplification system that is proved to be sound and complete, and (2) a proof showing that FT satisfies the so-called “independence of negative constraints”.

*Digital Equipment Corporation, Paris Research Laboratory (PRL), 85 avenue Victor Hugo, 92500 Rueil-Malmaison, France (email: {hak,podelski}@prl.dec.com).

†German Research Center for Artificial Intelligence (DFKI) and Universität des Saarlandes, Stuhlsatzenhausweg 3, 6600 Saarbrücken 11, Germany (email: smolka@dfki.uni-sb.de). Supported in part by the Bundesminister für Forschung und Technologie under contract ITW 9105.

Contents

1	Introduction	3
2	Feature Trees and Constraints	6
3	Basic Simplification	9
4	Entailment, Independence and Negation	12
5	Entailment Simplification	14
6	Conclusion	20

1 Introduction

An important structural property of many logic programming systems is the fact that they factorize into a constraint system and an extension facility. Colmerauer’s Prolog II [8] is an early language design making explicit use of this property. CLP (Constraint Logic Programming [10]), ALPS [16], CCP (Concurrent Constraint Programming [21]), and KAP (Kernel Andorra Prolog [9]) are recent logic programming frameworks that exploit this property to its full extent by being parameterized with respect to an abstract class of constraint systems. The basic operation these frameworks require of a constraint system is a test for unsatisfiability. ALPS, CCP, and KAP in addition require a test for entailment between constraints, which is needed for advanced control mechanisms such as delaying, coroutining, synchronisation, committed choice, and deep constraint propagation. Given this situation, constraint systems are a central issue in research on logic programming.

The constraint systems of most existing logic programming languages are variations and extensions of Herbrand [14], the constraint system underlying Prolog. The individuals of Herbrand are trees corresponding to ground terms, and the atomic constraints are equations between terms. Seen from the perspective of programming, Herbrand provides a universal data structure as a logical system.

This paper presents a constraint system FT , which we feel is an intriguing alternative to Herbrand both theoretically and practically. As does Herbrand, FT provides a universal data structure based on trees. However, the trees of FT (called feature trees) are more general than the trees of Herbrand (called constructor trees), and the constraints of FT are finer grained and of different expressivity. The basic notion of FT are functional attributes called features, which provide for record-like descriptions of data avoiding the overspecification intrinsic in Herbrand’s constructor-based descriptions. For the special case of constructor trees, features amount to argument selectors for constructors.

Suppose we want to say that x is a wine whose grape is **riesling** and whose color is **white**. To do this in Herbrand, one may write the equation

$$x = \text{wine}(\text{riesling}, \text{white}, y_1, \dots, y_n)$$

with the implicit assumption that the first argument of the constructor **wine** carries the “feature” **grape**, the second argument carries the “feature” **color**, and the remaining arguments y_1, \dots, y_n carry the remaining “features” of the chosen representation of wines. The obvious difficulty with this description is that it says more than we want to say, namely, that the constructor **wine** has $n + 2$ arguments and that the “features” **grape** and **color** are represented as the first and the second argument.




Figure 1: Examples of Feature Trees.

The constraint system FT avoids this overspecification by allowing the description

$$x: \text{wine}[\text{grape} \Rightarrow \text{riesling}, \text{color} \Rightarrow \text{white}] \quad (1)$$

saying that x has sort wine , its feature grape is riesling , and its feature color is white . Nothing is said about other features of x , which may or may not exist.

The individuals of FT are so-called feature trees, examples of which are shown in Figure 1. A feature tree is a possibly infinite tree whose nodes are labeled with symbols called sorts, and whose edges are labeled with symbols called features. The labeling with features is deterministic in that all edges departing from a node must be labeled with distinct features. Thus, every direct subtree of a feature tree can be identified by the feature labeling the edge leading to it. The constructor trees of Herbrand can be represented as feature trees whose edges are labeled with natural numbers indicating the corresponding argument positions.

All but the second and third feature tree in Figure 1 satisfy the description (1).

The constraints of FT are ordinary first-order formulae taken over a signature that accommodates sorts as unary and features as binary predicates. Thus the description (1) is actually syntactic sugar for the formula

$$\text{wine}(x) \wedge \exists y(\text{grape}(x, y) \wedge \text{riesling}(y)) \wedge \exists y(\text{color}(x, y) \wedge \text{white}(y)).$$

The set of all rational feature trees is made into a corresponding logical structure \mathcal{T} by letting $A(x)$ hold iff the root of x is labeled with the sort A , and letting $f(x, y)$ hold iff x has y as direct subtree via the feature f . The feature tree structure \mathcal{T} fixes an algebraic semantics for FT .

We will also establish a logical semantics, which is given by three axiom schemes fixing a first-order theory FT . Backofen and Smolka [6] show that \mathcal{T} is a model of FT and that FT is in fact a complete theory, which means that FT is exactly the theory induced by \mathcal{T} . However, we will not use the completeness result in the present paper, but show explicitly that entailment with respect to \mathcal{T} is the same as entailment with respect to FT .

The two major technical contributions of this paper are (1) an incremental entailment simplification system that is proved to be sound and complete, and (2) a proof showing that FT satisfies the so-called “independence of negative constraints” [7, 14, 15]. The incremental entailment simplification system is the prerequisite for FT ’s use with either of the constraint programming frameworks ALPS, CCP or KAP mentioned at the beginning of this section. The independence property means among other things that negative constraints can essentially be handled through entailment simplification.

One origin of FT is Ait-Kaci’s ψ -term calculus [1], which is at the heart of the programming language LOGIN [3] and further extended in the language LIFE [5] with functions over feature structures thanks to a generalization of the concept of residuation of Le Fun [4]. Other precursors of FT are the feature descriptions found in so-called unification grammars [13, 12] developed for natural language processing, and also the formalisms of Mukai [17, 18]. These early feature structure formalisms were presented in a nonlogical form. Major steps in the process of their understanding and logical reformulation are the articles [20, 23, 11, 22]. Feature trees, the feature tree structure \mathcal{T} , and the axiomatization of \mathcal{T} were first given in [6].

The paper is organized as follows. Section 2 defines the basic notions and discusses the differences in expressivity between Herbrand and FT . Section 3 gives a basic simplification system that decides satisfiability of positive constraints. Section 4 is not committed to FT but discusses the notion of incremental entailment checking and its connection with the independence property and negation. Section 5 gives the entailment simplification system, proves it sound, complete and terminating, and also proves that FT satisfies the independence property.

2 Feature Trees and Constraints

To give a rigorous formalization of feature trees, we first fix two disjoint alphabets \mathcal{S} and \mathcal{F} , whose symbols are called **sorts** and **features**, respectively. The letters A, B, C will always denote sorts, and the letters f, g, h will always denote features. Words over \mathcal{F} are called paths. The concatenation of two paths v and w results in the path vw . The symbol ε denotes the empty path, $v\varepsilon = \varepsilon v = v$, and \mathcal{F}^* denotes the set of all paths.

A **tree domain** is a nonempty set $D \subseteq \mathcal{F}^*$ that is prefix-closed, that is, if $vw \in D$, then $v \in D$. Thus, it always contains the empty path.

A **feature tree** is a mapping $t : D \rightarrow \mathcal{S}$ from a tree domain D into the set of sorts. The paths in the domain of a feature tree represent the nodes of the tree; the empty path represents its root. The letters s and t are used to denote feature trees.

If convenient, we consider a feature tree t as a relation, *i.e.*, $t \subseteq \mathcal{F}^* \times \mathcal{S}$, and write $(w, A) \in t$ instead of $t(w) = A$. As relations, *i.e.*, as subsets of $\mathcal{F}^* \times \mathcal{S}$, feature trees are partially ordered by set inclusion. We say that s is **smaller than** t if $s \subseteq t$.

The **subtree** wt of a feature tree t at one of its nodes w is the feature tree defined by (as a relation):

$$wt := \{(v, A) \mid (wv, A) \in t\}.$$

If D is the domain of t , then the domain of wt is the set $w^{-1}D = \{v \mid wv \in D\}$. Thus, wt is given as the mapping $wt : w^{-1}D \rightarrow \mathcal{S}$ defined on its domain by $wt(v) = t(wv)$. A feature tree s is called a **subtree** of a feature tree t if it is a subtree $s = wt$ at one of its nodes w , and a **direct subtree** if $w \in \mathcal{F}$.

A feature tree t with domain D is called **rational** if (1) t has only finitely many subtrees and (2) t is finitely branching, which is: for every $w \in D$, $w\mathcal{F} \cap D = \{wf \in D \mid f \in \mathcal{F}\}$ is finite. Assuming (1), (2) is equivalent to saying that there exist finitely many features f_1, \dots, f_n such that $D \subseteq \{f_1, \dots, f_n\}^*$.

Constraints over feature trees will be defined as first-order formulae. We first fix a first-order signature $\mathcal{S} \uplus \mathcal{F}$ by taking sorts as unary and features as binary relation symbols. Moreover, we fix an infinite alphabet of **variables** and adopt the convention that x, y, z always denote variables. Under this signature, every term is a variable and an **atomic formula** is either a feature constraint xfy ($f(x, y)$ in standard notation), a sort constraint Ax ($A(x)$ in standard notation), an equation $x \doteq y$, \perp (“false”), or \top (“true”). Compound formulae are obtained as usual by the connectives $\wedge, \vee, \rightarrow, \leftrightarrow, \neg$ and the quantifiers \exists and \forall . We use $\exists\phi$ and $\forall\phi$ to denote the existential

and universal closure of a formula ϕ , respectively. Moreover, $\mathcal{V}(\phi)$ is taken to denote the set of all variables that occur free in a formula ϕ . The letters ϕ and ψ will always denote formulae. In the following we won't make a distinction between formulae and constraints, that is, a **constraint** is a formula as defined above.

$\mathcal{S} \uplus \mathcal{F}$ -**structures** and **validity** of formulae in $\mathcal{S} \uplus \mathcal{F}$ -structures are defined as usual. Since we consider only $\mathcal{S} \uplus \mathcal{F}$ -structures in the following, we will simply speak of structures. A **theory** is a set of closed formulae. A **model** of a theory is a structure that satisfies every formulae of the theory. A formula ϕ is a **consequence** of a theory T ($T \models \phi$) if $\forall \phi$ is valid in every model of T . A formula ϕ is **satisfiable** in a structure \mathcal{A} if $\exists \phi$ is valid in \mathcal{A} . Two formulae ϕ, ψ are **equivalent** in a structure \mathcal{A} if $\forall (\phi \leftrightarrow \psi)$ is valid in \mathcal{A} . We say that a formula ϕ **entails** a formula ψ in a structure \mathcal{A} [theory T] and write $\phi \models_{\mathcal{A}} \psi$ [$\phi \models_T \psi$] if $\forall (\phi \rightarrow \psi)$ is valid in \mathcal{A} [is a consequence of T]. A theory T is **complete** if for every closed formula ϕ either ϕ or $\neg\phi$ is a consequence of T .

The **feature tree structure** \mathcal{T} is the $\mathcal{S} \uplus \mathcal{F}$ -structure defined as follows:

- the domain of \mathcal{T} is the set of all rational feature trees;
- $t \in A^{\mathcal{T}}$ iff $t(\varepsilon) = A$ (t 's root is labeled with A);
- $(s, t) \in f^{\mathcal{T}}$ iff $f \in D_s$ and $t = fs$ (t is the subtree of s at f).

Next we discuss the expressivity of our constraints with respect to feature trees (that is, with respect to the feature tree structure \mathcal{T}) by means of examples. The constraint

$$\neg \exists y(xfy)$$

says that x has no subtree at f , that is, that there is no edge departing from x 's root that is labeled with f . To say that x has subtree y at path $f_1 \cdots f_n$, we can use the constraint

$$\exists z_1 \cdots \exists z_{n-1}(xf_1z_1 \wedge z_1f_2z_2 \wedge \dots \wedge z_{n-1}f_ny).$$

Now let's look at statements we cannot express (more precisely, statements of whom the authors believe they cannot be expressed). One simple unexpressible statement is “ y is a subtree of x ” (that is, “ $\exists w: y = wx$ ”). Moreover, we cannot express that x is smaller than y . Finally, if we assume that the alphabet \mathcal{F} of features is infinite, we cannot say that x has subtrees at features f_1, \dots, f_n but no subtree at any other feature. In particular, we then cannot say that x is a primitive feature tree, that is, has no proper subtree.

The theory FT_0 is given by the following two axiom schemes:

(Ax1) $\forall x \forall y \forall z (xfy \wedge xfz \rightarrow y \doteq z)$
(for every feature f)

(Ax2) $\forall x (Ax \wedge Bx \rightarrow \perp)$
(for every two distinct sorts A and B).

The first axiom scheme says that features are functional and the second scheme says that sorts are mutually disjoint. Clearly, \mathcal{T} is a model of FT_0 . Moreover, FT_0 is incomplete (for instance, $\exists x(Ax)$ is valid in \mathcal{T} but invalid in other models of FT_0). We will see in the next section that FT_0 plays an important role with respect to basic constraint simplification.

Next we introduce some additional notation needed in the rest of the paper. This notation will also allow us to state a third axiom scheme that, as shown in [6], extends FT_0 to a complete axiomatization of \mathcal{T} .

Throughout the paper we assume that the conjunction of formulae is an associative and commutative operator that has \top as neutral element. This means that we identify $\phi \wedge (\psi \wedge \theta)$ with $\theta \wedge (\psi \wedge \phi)$, and $\phi \wedge \top$ with ϕ (but not, for example, $xfy \wedge xfy$ with xfy). A conjunction of atomic formulae can thus be seen as the finite multiset of these formulae, where conjunction is multiset union, and \top (the “empty conjunction”) is the empty multiset. We will write $\psi \subseteq \phi$ (or $\psi \in \phi$, if ψ is an atomic formula) if there exists a formula ψ' such that $\psi \wedge \psi' = \phi$.

We will use an additional atomic formula $xf\uparrow$ (“ f undefined on x ”) that is taken to be equivalent to $\neg\exists y(xfy)$, for some variable y (other than x).

Only for the formulation of the third axiom we introduce the notion of a **solved-clause**, which is either \top or a conjunction ϕ of atomic formulae of the form xfy , Ax or $xf\uparrow$ such that the following conditions are satisfied:

1. if $Ax \in \phi$ and $Bx \in \phi$, then $A = B$;
2. if $xfy \in \phi$ and $xfz \in \phi$, then $y = z$;
3. if $xfy \in \phi$, then $xf\uparrow \notin \phi$.

Given a solved-clause ϕ , we say that a variable x is **dependent** in ϕ if ϕ contains a constraint of the form Ax , xfy or $xf\uparrow$, and use $\mathcal{DV}(\phi)$ to denote the set of all variables that are dependent in ϕ .

The theory FT is obtained from FT_0 by adding the axiom scheme:

(Ax3) $\tilde{\forall} \exists X \phi$
(for every solved-clause ϕ and $X = \mathcal{DV}(\phi)$).

Theorem 2.1 *The feature tree structure \mathcal{T} is a model of the theory FT.*

Proof. We will only show that FT is a model of the third axiom. Let X be the set of dependent variables of the solved-clause ϕ , $X = \mathcal{DV}(\phi)$. Let α be any \mathcal{T} -valuation defined on $\mathcal{V}(\phi) - X$; we write the tree $\alpha(y)$ as t_y . We will extend α on X such that $\mathcal{T}, \alpha \models \phi$.

Given $x \in X$, we define the “punctual” tree $t_x = \{(\varepsilon, A)\}$, where $A \in \mathcal{S}$ is the sort such that $Ax \in \phi$, if it exists, and arbitrary, otherwise. Now we are going to use the notion of *tree sum* of Nivat [19], where $w^{-1}t = \{(wv, A) \mid (v, A) \in t\}$ (“the tree t translated by w ”), and we define:

$$\alpha(x) = \biguplus \{w^{-1}t_y \mid x \overset{w}{\rightsquigarrow} y \text{ for some } y \in \mathcal{V}(\phi), w \in \mathcal{F}^*\}.$$

Here the “leads-to” relation $\overset{w}{\rightsquigarrow}$ is given by: $x \overset{\varepsilon}{\rightsquigarrow} x$, and $x \overset{wf}{\rightsquigarrow} y$ if $x \overset{w}{\rightsquigarrow} y'$ and $y'fy \in \phi$, for some $y' \in \mathcal{V}(\phi)$ and some $f \in \mathcal{F}$. Since

$$\alpha(x) = \bigcup \{w^{-1}\alpha(y) \mid \dots\}$$

and $w\alpha(x) = \alpha(y)$, it follows that $\alpha(x)$ is a rational tree and that $\mathcal{T}, \alpha \models \phi$. □

3 Basic Simplification

A **basic constraint** is either \perp or a possibly empty conjunction of atomic formulae of the form Ax , xfy , and $x \doteq y$. The following five **basic simplification** rules constitute a simplification system for basic constraints, which, as we will see, decides whether a basic constraint is satisfiable in \mathcal{T} .

1.
$$\frac{xfy \wedge x fz \wedge \phi}{xfz \wedge y \doteq z \wedge \phi}$$
2.
$$\frac{Ax \wedge Bx \wedge \phi}{\perp} \quad A \neq B$$
3.
$$\frac{Ax \wedge Ax \wedge \phi}{Ax \wedge \phi}$$
4.
$$\frac{x \doteq y \wedge \phi}{x \doteq y \wedge \phi[x \leftarrow y]} \quad x \in \mathcal{V}(\phi) \text{ and } x \neq y$$

$$5. \frac{x \doteq x \wedge \phi}{\phi}$$

The notation $\phi[x \leftarrow y]$ is used to denote the formula that is obtained from ϕ by replacing every occurrence of x with y . We say that a constraint ϕ **simplifies to** a constraint ψ by a simplification rule ρ if $\frac{\phi}{\psi}$ is an instance of ρ . We say that a constraint ϕ **simplifies to** a constraint ψ if either $\phi = \psi$ or ϕ simplifies to ψ in finitely many steps each licensed by one of the five simplification rules given above.

Example 3.1 We have the following basic simplification chain, leading to a solved constraint:

$$\begin{aligned} & xfu \wedge yfv \wedge Au \wedge Av \wedge z \doteq x \wedge y \doteq z \\ \Rightarrow & xfu \wedge yfv \wedge Au \wedge Av \wedge z \doteq x \wedge y \doteq x \\ \Rightarrow & xfu \wedge xfv \wedge Au \wedge Av \wedge z \doteq x \wedge y \doteq x \\ \Rightarrow & xfv \wedge Au \wedge Av \wedge u \doteq v \wedge z \doteq x \wedge y \doteq x \\ \Rightarrow & xfv \wedge Av \wedge Av \wedge u \doteq v \wedge z \doteq x \wedge y \doteq x \\ \Rightarrow & xfv \wedge Av \wedge u \doteq v \wedge z \doteq x \wedge y \doteq x \end{aligned}$$

Using the same steps up to the last one, the constraint $xfu \wedge yfv \wedge Au \wedge Bv \wedge z \doteq x \wedge y \doteq z$ simplifies to \perp (in the last step, Rule 2 instead of Rule 3 is applied). \square

Proposition 3.2 *If the basic constraint ϕ simplifies to ψ , then $FT_0 \models \phi \leftrightarrow \psi$.*

Proof. The rules 3, 4 and 5 perform equivalence transformations with respect to every structure. The rules 1 and 2 correspond exactly to the two axiom schemes of FT_0 and perform equivalence transformations with respect to every model of FT_0 . \square

We say that a basic constraint ϕ **binds** a variable x to y if $x \doteq y \in \phi$ and x occurs only once in ϕ . At this point it is important to note that we consider equations as ordered, that is, assume that $x \doteq y$ is different from $y \doteq x$ if $x \neq y$. We say that a variable x is **eliminated**, or **bound by** ϕ , if ϕ binds x to some variable y .

Proposition 3.3 *The basic simplification rules are terminating.*

Proof. First observe that the simplification rules don't add new variables and preserve eliminated variables. Furthermore, rule 4 increases the number of eliminated variables by one. Hence we know that if an infinite simplification chain exists, we can assume without loss of generality that it only employs the rules 1, 3 and 5. Since rule 1 decreases the number of feature constraints “ xfy ”, which is not increased by rules 3 and 5, we know that if an infinite simplification chain exists, we can assume without loss of generality that it only employs the rules 3 and 5. Since this is clearly impossible, an infinite simplification chain cannot exist. \square

A basic constraint is called **normal** if none of the five simplification rules applies to it. A constraint ψ is called a **normal form** of a basic constraint ϕ if ϕ can be simplified to ψ and ψ is normal. A **solved constraint** is a normal constraint that is different from \perp .

So far we know that we can compute for any basic constraint ϕ a normal form ψ by applying the simplification rules as long as they are applicable. Although the normal form ψ may not be unique for ϕ , we know that ϕ and ψ are equivalent in every model of FT_0 . It remains to show that every solved constraint is satisfiable in \mathcal{T} .

Every basic constraint ϕ has a unique decomposition $\phi = \phi_N \wedge \phi_G$ such that ϕ_N is a possibly empty conjunction of equations “ $x \doteq y$ ” and ϕ_G is a possibly empty conjunction of feature constraints “ xfy ” and sort constraints “ Ax ”. We call ϕ_N the **normalizer** and ϕ_G the **graph** of ϕ .

Proposition 3.4 *A basic constraint $\phi \neq \perp$ is solved iff the following conditions hold:*

1. *an equation $x \doteq y$ appears in ϕ only if x is eliminated in ϕ ;*
2. *the graph of ϕ is a solved clause;*
3. *no primitive constraint appears more than once in ϕ .*

Proposition 3.5 *Every solved constraint is satisfiable in every model of FT .*

Proof. Let ϕ be a solved constraint and \mathcal{A} be a model of FT . Then we know by axiom scheme Ax3 that the graph ϕ_G of a solved constraint ϕ is satisfiable in an FT -model \mathcal{A} . A variable valuation α into \mathcal{A} such that $\mathcal{A}, \alpha \models \phi_G$ can be extended on all eliminated variables simply by $\alpha(x) = \alpha(y)$ if $x \doteq y \in \phi$, such that $\mathcal{A}, \alpha \models \phi$. \square

Theorem 3.6 *Let ψ be a normal form of a basic constraint ϕ . Then ϕ is satisfiable in \mathcal{T} if and only if $\psi \neq \perp$.*

Proof. Since ϕ and ψ are equivalent in every model of FT_0 and \mathcal{T} is a model of FT_0 , it suffices to show that ψ is satisfiable in \mathcal{T} if and only if $\psi \neq \perp$. To show the nontrivial direction, suppose $\psi \neq \perp$. Then ψ is solved and we know by the preceding proposition that ψ is satisfiable in every model of FT . Since \mathcal{T} is a model of FT , we know that ψ is satisfiable in \mathcal{T} . \square

Theorem 3.7 *For every basic constraint ϕ the following statements are equivalent:*

$$\mathcal{T} \models \tilde{\exists}\phi \Leftrightarrow \exists \text{ model } \mathcal{A} \text{ of } FT_0: \mathcal{A} \models \tilde{\exists}\phi \Leftrightarrow FT \models \tilde{\exists}\phi.$$

Proof. The implication $1 \Rightarrow 2$ holds since \mathcal{T} is a model of FT_0 . The implication $3 \Rightarrow 1$ follows from the fact that \mathcal{T} is a model of FT . It remains to show that $2 \Rightarrow 3$.

Let ϕ be satisfiable in some model of FT_0 . Then we can apply the simplification rules to ϕ and compute a normal form ψ such that ϕ and ψ are equivalent in every model of FT_0 . Hence ψ is satisfiable in some model of FT_0 . Thus $\psi \neq \perp$, which means that ψ is solved. Hence we know by the preceding proposition that ψ is satisfiable in every model of FT . Since ϕ and ψ are equivalent in every model of $FT_0 \subseteq FT$, we have that ϕ is satisfiable in every model of FT . \square

4 Entailment, Independence and Negation

In this section we discuss some general properties of constraint entailment. This prepares the ground for the next section, which is concerned with entailment simplification in the feature tree constraint system.

Throughout this section we assume that \mathcal{A} is a structure, γ and ϕ are formulae that can be interpreted in \mathcal{A} , and that X is a finite set of variables.

We say that γ **disentails** ϕ in \mathcal{A} if γ entails $\neg\phi$ in \mathcal{A} . If γ is satisfiable in \mathcal{A} , then γ cannot both entail and disentail $\exists X\phi$ in \mathcal{A} . We say that γ **determines** ϕ in \mathcal{A} if γ either entails or disentails ϕ in \mathcal{A} .

Given γ , ϕ and X , we want to determine in an **incremental** manner whether γ entails or disentails $\exists X\phi$. Typically, γ will **not** determine $\exists X\phi$ when $\exists X\phi$ is considered first, but this may change when γ is strengthened to $\gamma \wedge \gamma'$. The basic idea leading to an incremental entailment checker is to simplify ϕ

with respect to the **context** γ and the **local variables** X . Given γ , X and ϕ , simplification must yield a formula ψ such that

$$\gamma \models_{\mathcal{A}} \exists X \phi \leftrightarrow \exists X \psi.$$

The following facts provide some evidence that this is the right invariant for entailment simplification.

Proposition 4.1 *Let $\gamma \models_{\mathcal{A}} \exists X \phi \leftrightarrow \exists X \psi$. Then:*

1. $\gamma \models_{\mathcal{A}} \exists X \phi$ iff $\gamma \models_{\mathcal{A}} \exists X \psi$;
2. $\gamma \models_{\mathcal{A}} \neg \exists X \phi$ iff $\gamma \models_{\mathcal{A}} \neg \exists X \psi$;
3. if $\psi = \perp$, then $\gamma \models_{\mathcal{A}} \neg \exists X \phi$;
4. if $\exists X \psi$ is valid in \mathcal{A} , then $\gamma \models_{\mathcal{A}} \exists X \phi$.

Statements 1 and 2 say that it doesn't matter whether entailment and dis-entailment are decided for ϕ or ψ . Statement 3 gives a local condition for disentanglement, and Statement 4 gives a local condition for entailment. The entailment simplification system for feature trees given in the next section will in fact decide entailment and disentanglement by simplifying such that the condition of Statement 4 is met in the case of entailment, and that the condition of Statement 3 is met in the case of disentanglement.

In practice, one can ensure by variable renaming that no variable of X occurs in γ . The next fact says that then it suffices if entailment simplification respects the more convenient invariant

$$\mathcal{A} \models \gamma \wedge \phi \leftrightarrow \gamma \wedge \psi.$$

This is the invariant respected by our system (**cf.** Proposition 5.4).

Proposition 4.2 *Let $X \cap \mathcal{V}(\gamma) = \emptyset$. Then:*

1. if $\mathcal{A} \models \gamma \wedge \phi \leftrightarrow \gamma \wedge \psi$, then $\gamma \models_{\mathcal{A}} \exists X \phi \leftrightarrow \exists X \psi$;
2. $\gamma \models_{\mathcal{A}} \neg \exists X \phi$ iff $\gamma \wedge \phi$ is unsatisfiable in \mathcal{A} .

That is, the conjunction $\gamma \wedge \phi$ is satisfiable if and only if γ either entails $\exists X \phi$, or it does not determine $\exists X \phi$.

The so-called independence of negative constraints [7, 14, 15] is an important property of constraint systems. If it holds, simplification of conjunctions of

positive and negative constraints can be reduced to entailment simplification of conjunctions of positive constraints.

To define the independence property, we assume that a constraint system is a pair consisting of a structure \mathcal{A} and a set of so-called basic constraints. From basic constraints one can build more complex constraints using the connectives and quantifiers of predicate logic. We say that a constraint system satisfies the **independence property** if

$$\gamma \models_{\mathcal{A}} \exists X_1 \phi_1 \vee \dots \vee \exists X_n \phi_n \text{ iff } \exists i: \gamma \models_{\mathcal{A}} \exists X_i \phi_i$$

for all basic constraints $\gamma, \phi_1, \dots, \phi_n$ and all finite sets of variables X_1, \dots, X_n .

Proposition 4.3 *If a constraint system satisfies the independence property, then the following statements hold (γ, ϕ and ϕ_1, \dots, ϕ_n are basic constraints):*

1. $\gamma \wedge \neg \exists X_1 \phi_1 \wedge \dots \wedge \neg \exists X_n \phi_n$ *unsatisfiable in \mathcal{A} iff $\exists i: \gamma \models_{\mathcal{A}} \exists X_i \phi_i$;*
2. *if $\gamma \wedge \neg \exists X_1 \phi_1 \wedge \dots \wedge \neg \exists X_n \phi_n$ is satisfiable in \mathcal{A} , then $\gamma \wedge \neg \exists X_1 \phi_1 \wedge \dots \wedge \neg \exists X_n \phi_n \models_{\mathcal{A}} \exists X \phi$ iff $\gamma \models_{\mathcal{A}} \exists X \phi$.*

5 Entailment Simplification

We now return to the feature tree constraint system. Throughout this section we assume that γ is a solved constraint and X is a finite set of variables not occurring in γ . We will call γ the **context**, the variables in X **local**, and all other variables **global**.

If T is a theory and ϕ and ψ are possibly open formulae, we write $\phi \models_T \psi$ (read: ϕ entails ψ in T) if $\forall (\phi \rightarrow \psi)$ is valid in T .

Theorem 5.1 *For every basic constraint ϕ , the following equivalences hold:*

$$\gamma \models_{\mathcal{T}} \neg \exists X \phi \text{ iff } \gamma \models_{FT_0} \neg \exists X \phi \text{ iff } \gamma \models_{FT} \neg \exists X \phi.$$

Proof. Implication “2 \Rightarrow 3” holds since $FT_0 \subseteq FT$. Implication “3 \Rightarrow 1” holds since \mathcal{T} is a model of FT . To show implication “1 \Rightarrow 2”, suppose $\gamma \models_{\mathcal{T}} \neg \exists X \phi$. Then we know by Proposition 4.2 that $\gamma \wedge \phi$ is unsatisfiable in \mathcal{T} . Thus we know by Theorem 3.7 that $\gamma \wedge \phi$ is unsatisfiable in every model of FT_0 . Hence we know by Proposition 4.2 that $\gamma \models_{FT_0} \neg \exists X \phi$. \square

For every basic constraint ϕ and every variable x we define

$$\phi x := \begin{cases} y & \text{if } x \doteq y \in \phi \text{ and } x \text{ is eliminated;} \\ x & \text{otherwise.} \end{cases}$$

A basic constraint ϕ is **X -oriented** if $x \doteq y \in \phi$ always implies $x \in X$ or $y \notin X$. A basic constraint ϕ is **pivoted** if $x \doteq y \in \phi$ implies that x is eliminated in ϕ (and then y is a “pivot”).

The following **entailment simplification rules** simplify basic constraints to basic constraints with respect to a context γ and local variables X .

1. $\frac{xfu \wedge \phi}{u \doteq v \wedge \phi} \quad yfv \in \gamma \wedge \phi, \quad \phi y = x$
2. $\frac{\phi}{\phi u \doteq \phi v \wedge \phi} \quad \begin{cases} xfu \wedge yfv \subseteq \gamma, \\ \phi x = \phi y, \quad \phi u \neq \phi v, \\ \phi \text{ } X\text{-oriented and pivoted} \end{cases}$
3. $\frac{\phi}{\perp} \quad Ax \wedge By \subseteq \gamma \wedge \phi, \quad \phi x = \phi y, \quad A \neq B$
4. $\frac{Ax \wedge \phi}{\phi} \quad Ay \in \gamma \wedge \phi, \quad \phi y = x$
5. $\frac{x \doteq y \wedge \phi}{x \doteq y \wedge \phi[x \leftarrow y]} \quad \begin{cases} x \neq y, \quad x \in \mathcal{V}(\phi), \\ (x \in X \text{ or } y \notin X) \end{cases}$
6. $\frac{x \doteq y \wedge \phi}{y \doteq x \wedge \phi} \quad x \notin X, \quad y \in X$
7. $\frac{\phi}{\phi[x \leftarrow y]} \quad x \doteq y \in \gamma, \quad x \in \mathcal{V}(\phi)$
8. $\frac{x \doteq x \wedge \phi}{\phi}$

We say that a basic constraint ϕ **simplifies** to a constraint ψ **with respect to γ and X** if $\phi = \psi$ or ϕ simplifies to ψ in finitely many steps each licensed by one of the eight simplification rules given above. The notions of **normal** and **normal form with respect to γ** are defined accordingly.

Example 5.2 Let $\gamma = xfu \wedge yfv \wedge Au \wedge Bv$ and $X = \{z\}$. Then we have the following simplification chain with respect to γ and X :

$$\begin{array}{lll}
& x \doteq z \wedge y \doteq z & \\
\Rightarrow_{\gamma, X} & z \doteq x \wedge y \doteq z & \text{by Rule E6} \\
\Rightarrow_{\gamma, X} & z \doteq x \wedge y \doteq x & \text{by Rule E5} \\
\Rightarrow_{\gamma, X} & u \doteq v \wedge z \doteq x \wedge y \doteq x & \text{by Rule E2} \\
\Rightarrow_{\gamma, X} & \perp & \text{by Rule E3.}
\end{array}$$

Let us now take as context $\tilde{\gamma} = xfu \wedge yfv \wedge Au$. Then $\tilde{\phi} = u \doteq v \wedge z \doteq x \wedge y \doteq x$ is normal with respect to $\tilde{\gamma}$ and X . We shall see that this normal form tells us that $\tilde{\gamma}$ does not determine $\tilde{\phi}$. If $\tilde{\gamma}$ gets strengthened either to $\tilde{\gamma} \wedge Bv$ (as above), or to $\tilde{\gamma} \wedge x \doteq y$, then the strengthened context does determine: it disentails in the first and entails in the second case. The basic normal form of $\tilde{\gamma} \wedge x \doteq y$ is $yfu \wedge Au \wedge v \doteq u \wedge x \doteq y$; with respect to this context $\tilde{\phi}$ simplifies to $z \doteq y$. \square

In the previous example, $\phi = z \doteq x \wedge y \doteq x$ simplifies to $\phi_1 = u \doteq v \wedge z \doteq x \wedge y \doteq x$ with respect to $\gamma = xfu \wedge yfv \wedge Au \wedge Bv$ and $X = \{z\}$. This corresponds to a basic simplification as follows:

$$\begin{array}{l}
\gamma \wedge \phi = \\
\Rightarrow xfu \wedge yfv \wedge Au \wedge Bv \quad \wedge z \doteq x \wedge y \doteq x \\
\Rightarrow xfu \wedge xfv \wedge Au \wedge Bv \quad \wedge z \doteq x \wedge y \doteq x \\
\Rightarrow xfv \wedge Au \wedge Bv \quad \wedge u \doteq v \wedge z \doteq x \wedge y \doteq x \\
= \gamma' \wedge \phi'_1
\end{array}$$

We observe that $\gamma \wedge \phi_1$ is equal to $\gamma' \wedge \phi'_1$, modulo renaming y by $\phi_1 y = x$ and u by $\phi_1 u = v$, and modulo the repetition of xfv .

Lemma 5.3 *Let ϕ simplify to ϕ_1 with respect to γ and X , not using Rule E6 (in an entailment simplification step). Then $\gamma \wedge \phi$ simplifies to some $\gamma' \wedge \phi'_1$ which is equal to $\gamma \wedge \phi_1$ up to variable renaming and repetition of conjuncts.*

Proof. Clearly, each entailment simplification rule, except for E6, corresponds directly to a basic simplification rule (namely, E1 and E2 to B1, E3 to B2, E4 to B3, E5 and E7 to B4, and E8 to B5).

If the application of the entailment simplification rule to ϕ relies on a condition of the form $\phi x = y$ or $\phi x = \phi y$ where $x \neq \phi x$ or $y \neq \phi y$, then $x \doteq \phi x \in \phi$ or $y \doteq \phi y \in \phi$, and Rule B4 is first applied to $\gamma \wedge \phi$, eliminating x by ϕx (y by ϕy).

When comparing $\gamma \wedge \phi_1$ and $\gamma' \wedge \phi'_1$, renamings take account of these variable eliminations. Note that, if the rule applied to ϕ is E2, then γ' has one feature constraint xfv less than γ — which, after renaming, has a repetition of exactly this constraint. \square

Proposition 5.4 *If ϕ simplifies to ψ with respect to γ and X , then $\gamma \wedge \phi$ and $\gamma \wedge \psi$ are equivalent in every model of FT_0 .*

Proof. Follows from Lemma 5.3 and Proposition 3.2. \square

Proposition 5.5 *The entailment simplification rules are terminating, provided γ and X are fixed.*

Proof. First we strengthen the statement by weakening the applicability conditions $\phi y = x$ in Rules E1 and E4 to $\phi y = \phi x$. Then from Lemma 5.3 follows: (*) Each entailment simplification rule applies to ϕ_1 with respect to γ and X if and only if it applies to ϕ'_1 with respect to γ' and X — except possibly for E5, when the corresponding variable has already been eliminated in an “extra” basic simplification step.

If γ' has one conjunct of the form xfu less than γ , then (*) still holds; regarding a new application of E2 this is ensured by its (therefore so complicated...) applicability condition.

With condition (*), it is possible to prove by induction on n : For every entailment simplification chain $\phi, \phi_1, \dots, \phi_n$ with respect to γ and X , there exists a ‘basic plus Rule E6’ simplification chain $\gamma \wedge \phi, \gamma_1 \wedge \phi'_1, \dots, \gamma_{n+k} \wedge \phi'_{n+k}$, where $k \geq 0$ is the number of “extra” variable elimination steps. Since, according to Proposition 3.3, basic simplification chains are finite, so are entailment simplification chains. \square

So far we know that we can compute for any basic constraint ϕ a normal form ψ with respect to γ and X by applying the simplification rules as long as they are applicable. Although the normal form ψ may not be unique, we know that $\gamma \wedge \phi$ and $\gamma \wedge \psi$ are equivalent in every model of FT_0 .

Proposition 5.6 *For every basic constraint ϕ one can compute a normal form ψ with respect to γ and X . Every such normal form ψ satisfies: $\gamma \models_{\mathcal{T}} \exists X \phi$ iff $\gamma \models_{\mathcal{T}} \exists X \psi$, and $\gamma \models_{FT} \exists X \phi$ iff $\gamma \models_{FT} \exists X \psi$.*

Proof. Follows from Propositions 5.4, 5.5, 4.2 and 4.1. \square

In the following we will show that from the entailment normal form ψ of ϕ with respect to γ it is easy to tell whether we have entailment, disentanglement or neither. Moreover, the basic normal form of $\gamma \wedge \phi$ is exactly $\gamma \wedge \psi$ in the first case (and in the second, where $\gamma \wedge \perp = \perp$), and “almost” in the third case (**cf.** Lemma 5.3).

Proposition 5.7 *A basic constraint $\phi \neq \perp$ is normal with respect to γ and X if and only if the following conditions are satisfied:*

1. ϕ is solved, X -oriented, and contains no variable that is bound by γ ;
2. if $\phi x = y$ and $x f u \in \gamma$, then $y f v \notin \phi$ for every v ;
3. if $\phi x = \phi y$ and $x f u \in \gamma$ and $y f v \in \gamma$, then $\phi u = \phi v$;
4. if $\phi x = y$ and $A x \in \gamma$, then $B y \notin \phi$ for every B ;
5. if $\phi x = \phi y$ and $A x \in \gamma$ and $B y \in \gamma$, then $A = B$.

Lemma 5.8 *If $\phi \neq \perp$ is normal with respect to γ and X , then $\gamma \wedge \phi$ is satisfiable in every model of FT .*

Proof. Let $\phi \neq \perp$ be normal with respect to γ and X . Furthermore, let $\gamma = \gamma_N \wedge \gamma_G$ and $\phi = \phi_N \wedge \phi_G$ be the unique decompositions in normalizer and graph. Since the variables bound by γ_N occur neither in γ_G nor in ϕ , it suffices to show that $\gamma_G \wedge \phi_N \wedge \phi_G$ is satisfiable in every model of FT .

Let $\phi_N(\gamma_G)$ be the basic constraint that is obtained from γ_G by applying all bindings of ϕ_N . Then $\gamma_G \wedge \phi_N \wedge \phi_G$ is equivalent to $\phi_N \wedge \phi_N(\gamma_G) \wedge \phi_G$ and no variable bound by ϕ_N occurs in $\phi_N(\gamma_G) \wedge \phi_G$. Hence it suffices to show that $\phi_N(\gamma_G) \wedge \phi_G$ is satisfiable in every model of FT . With the conditions 2–5 of the preceding proposition it is easy to see that $\phi_N(\gamma_G) \wedge \phi_G$ is a solved clause. Hence we know by axiom scheme Ax3 that $\phi_N(\gamma_G) \wedge \phi_G$ is satisfiable in every model of FT . \square

Theorem 5.9 (Disentailment) *Let ψ be a normal form of ϕ with respect to γ and X . Then $\gamma \models_{\mathcal{T}} \neg \exists X \phi$ iff $\psi = \perp$.*

Proof. Suppose $\psi = \perp$. Then $\gamma \models_{\mathcal{T}} \neg \exists X \psi$ and hence $\gamma \models_{\mathcal{T}} \neg \exists X \phi$ by Proposition 5.6.

To show the other direction, suppose $\gamma \models_{\mathcal{T}} \neg \exists X \phi$. Then $\gamma \models_{\mathcal{T}} \neg \exists X \psi$ by Proposition 5.6 and hence $\gamma \wedge \psi$ unsatisfiable in \mathcal{T} by Proposition 4.2. Since \mathcal{T} is a model of FT (Theorem 2.1), we know by the preceding lemma that $\psi = \perp$ (since ψ is assumed to be normal). \square

We say that a variable x is **dependent** in a solved constraint ϕ if ϕ contains a constraint of the form Ax , xfy or $x \doteq y$. (Recall that equations are ordered;

thus y is not dependent in the constraint $x \doteq y$.) We use $\mathcal{DV}(\phi)$ to denote the set of all variables that are dependent in a solved constraint ϕ .

In the following we will assume that the underlying signature $\mathcal{S} \uplus \mathcal{F}$ has at least one sort and at least one feature that does not occur in the constraints under consideration. This assumption is certainly satisfied if the signature has infinitely many sorts and infinitely many features.

Lemma 5.10 (Spiting) *Let ϕ_1, \dots, ϕ_n be basic constraints different from \perp , and X_1, \dots, X_n be finite sets of variables disjoint from $\mathcal{V}(\gamma)$. Moreover, for every $i = 1, \dots, n$, let ϕ_i be normal with respect to γ and X_i , and let ϕ_i have a dependent variable that is not in X_i . Then $\gamma \wedge \neg \exists X_1 \phi_1 \wedge \dots \wedge \neg \exists X_n \phi_n$ is satisfiable in every model of FT .*

Proof. Let $\gamma = \gamma_N \wedge \gamma_G$ be the unique decomposition of γ into normalizer and graph. Since the variables bound by γ_N occur neither in γ_G nor in any ϕ_i , it suffices to show that $\gamma_G \wedge \neg \exists X_1 \phi_1 \wedge \dots \wedge \neg \exists X_n \phi_n$ is satisfiable in every model of FT . Thus it suffices to exhibit a solved clause δ such that $\gamma_G \subseteq \delta$ and, for every $i = 1, \dots, n$, $\mathcal{V}(\delta)$ is disjoint with X_i and $\delta \wedge \phi_i$ is unsatisfiable in every model of FT .

Without loss of generality we can assume that every X_i is disjoint with $\mathcal{V}(\gamma)$ and $\mathcal{V}(\phi_j) - X_j$ for all j . Hence we can pick in every ϕ_i a dependent variable x_i such that $x_i \notin X_j$ for any j .

Let z_1, \dots, z_k be all variables that occur on either side of equation $x_i \doteq y \in \phi_i$, $i = 1, \dots, n$ (recall that x_i is fixed for i). None of these variables occurs in any X_j since every ϕ_i is X_i -oriented. Next we fix a feature g and a sort B such that neither occurs in γ or any ϕ_i .

Now δ is obtained from γ by adding constraints as follows: if $Ax_i \in \phi_i$, then add Bx_i ; if $x_i f y \in \phi_i$, then add $x_i f \uparrow$; to enforce that the variables z_1, \dots, z_k are pairwise distinct, add

$$z_k g z_{k-1} \wedge \dots \wedge z_2 g z_1 \wedge z_1 g \uparrow .$$

It is straightforward to verify that these additions to γ yield a solved clause δ as required. \square

Proposition 5.11 *If ϕ is solved and $\mathcal{DV}(\phi) \subseteq X$, then $FT \models \tilde{\forall} \exists X \phi$.*

Proof. Let $\phi = \phi_N \wedge \phi_G$ be the decomposition of ϕ in normalizer and graph. Since every variable bound by ϕ is in X , it suffices to show that $\tilde{\forall} \exists X \phi_G$ is a consequence of FT . This follows immediately from axiom scheme $Ax3$ since ϕ_G is a solved clause. \square

Theorem 5.12 (Entailment) *Let ψ be a normal form of ϕ with respect to γ and X . Then $\gamma \models_{\mathcal{T}} \exists X\phi$ iff $\psi \neq \perp$ and $\mathcal{DV}(\psi) \subseteq X$.*

Proof. Suppose $\gamma \models_{\mathcal{T}} \exists X\phi$. Then we know $\gamma \models_{\mathcal{T}} \exists X\psi$ by Proposition 5.6, and thus $\gamma \wedge \neg\exists X\psi$ is unsatisfiable in \mathcal{T} . Since γ is solved, we know that γ is satisfiable in \mathcal{T} and hence that $\gamma \wedge \exists X\psi$ is satisfiable in \mathcal{T} . Thus $\psi \neq \perp$. Since $\gamma \wedge \neg\exists X\psi$ is unsatisfiable in \mathcal{T} and \mathcal{T} is a model of FT , we know by Lemma 5.10 that $\mathcal{DV}(\psi) \subseteq X$.

To show the other direction, suppose $\psi \neq \perp$ and $\mathcal{DV}(\psi) \subseteq X$. Then $FT \models \check{\forall}\exists X\psi$ by Proposition 5.11, and hence $\mathcal{T} \models \check{\forall}\exists X\psi$. Thus $\gamma \models_{\mathcal{T}} \exists X\psi$, and hence $\gamma \models_{\mathcal{T}} \exists X\phi$ by Proposition 5.6. \square

Theorem 5.13 *Let ϕ be a basic constraint. Then $\gamma \models_{\mathcal{T}} \exists X\phi$ iff $\gamma \models_{FT} \exists X\phi$.*

Proof. One direction holds since \mathcal{T} is a model of FT . To show the other direction, suppose $\gamma \models_{\mathcal{T}} \exists X\phi$. Without loss of generality we can assume that ϕ is normal with respect to γ and X . Hence we know by Theorem 5.12 that $\phi \neq \perp$ and $\mathcal{DV}(\phi) \subseteq X$. Thus $FT \models \check{\forall}\exists X\phi$ by Proposition 5.11 and hence $\gamma \models_{FT} \exists X\phi$. \square

Theorem 5.14 (Independence) *Let ϕ_1, \dots, ϕ_n be basic constraints, and X_1, \dots, X_n be finite sets of variables. Then*

$$\gamma \models_{\mathcal{T}} \exists X_1\phi_1 \vee \dots \vee \exists X_n\phi_n \text{ iff } \exists i: \gamma \models_{\mathcal{T}} \exists X_i\phi_i.$$

Proof. To show the nontrivial direction, suppose $\gamma \models_{\mathcal{T}} \exists X_1\phi_1 \vee \dots \vee \exists X_n\phi_n$. Without loss of generality we can assume that, for all $i = 1, \dots, n$, X_i is disjoint from $\mathcal{V}(\gamma)$, ϕ_i is normal with respect to γ and X_i , and $\phi_i \neq \perp$. Since $\gamma \wedge \neg\exists X_1\phi_1 \wedge \dots \wedge \neg\exists X_n\phi_n$ is unsatisfiable in \mathcal{T} and \mathcal{T} is a model of FT , we know by Lemma 5.10 that $\mathcal{DV}(\phi_k) \subseteq X_k$ for some k . Hence $\gamma \models_{\mathcal{T}} \exists X_k\phi_k$ by Theorem 5.12. \square

6 Conclusion

We have presented a constraint system FT for logic programming providing a universal data structure based on rational feature trees. FT accommodates record-like descriptions, which we think are superior to the constructor-based descriptions of Herbrand.

The declarative semantics of FT is specified both algebraically (the feature tree structure \mathcal{T}) and logically (the first-order theory FT given by three axiom schemes).

The operational semantics for FT is given by an incremental constraint simplification system, which can check satisfiability of and entailment between constraints. Since FT satisfies the independence property, the simplification system can also check satisfiability of conjunctions of positive and negative constraints.

We see four directions for further research.

First, FT should be strengthened such that it subsumes the expressivity of rational constructor trees [7, 8]. As is, FT cannot express that x is a tree having direct subtrees at exactly the features f_1, \dots, f_n . It turns out that the system CFT [24] obtained from FT by adding the primitive constraint

$$x\{f_1, \dots, f_n\}$$

(x has direct subtrees at exactly the features f_1, \dots, f_n) has the same nice properties as FT . In contrast to FT , CFT can express constructor constraints; for instance, the constructor constraint $x \doteq A(y, z)$ can be expressed equivalently as $Ax \wedge x\{1, 2\} \wedge x1y \wedge x2z$, if we assume that A is a sort and the numbers 1, 2 are features.

Second, it seems attractive to extend FT such that it can accommodate a sort lattice as used in [1, 3, 4, 5, 23]. One possibility to do this is to assume a partial order \leq on sorts and replace sort constraints Ax with *quasi-sort constraints* $[A]x$ whose declarative semantics is given as

$$[A]x \equiv \bigvee_{B \leq A} Bx.$$

Given the assumption that the sort ordering \leq has greatest lower bounds if lower bounds exist, it seems that the results and the simplification system given for FT carry over with minor changes.

Third, the worst-case complexity of entailment checking in FT should be established. We conjecture it to be quasi-linear in the size of γ and ϕ , provided the available features are fixed a priori.

Fourth, implementation techniques for FT at the level of the Warren abstract machine [2] need to be developed.

References

- [1] H. Ait-Kaci. An algebraic semantics approach to the effective resolution of type equations. *Theoretical Computer Science*, 45:293–351, 1986.

- [2] H. Aït-Kaci. *Warren's Abstract Machine: A Tutorial Reconstruction*. The MIT Press, Cambridge, MA, 1991.
- [3] H. Aït-Kaci and R. Nasr. LOGIN: A logic programming language with built-in inheritance. *The Journal of Logic Programming*, 3:185–215, 1986.
- [4] H. Aït-Kaci and R. Nasr. Integrating logic and functional programming. *Lisp and Symbolic Computation*, 2:51–89, 1989.
- [5] H. Aït-Kaci and A. Podelski. Towards a Meaning of LIFE. *Proceedings of the 3rd International Symposium on Programming Language Implementation and Logic Programming (Passau, Germany)*, J. Maluszyński and M. Wirsing, editors. LNCS 528, pages 255–274, Springer-Verlag, 1991.
- [6] R. Backofen and G. Smolka. A complete and decidable feature theory. Draft, German Research Center for Artificial Intelligence (DFKI), Stuhlsatzenhausweg 3, 6600 Saarbrücken 11, Germany, 1991. To appear.
- [7] A. Colmerauer. Equations and inequations on finite and infinite trees. In *Proceedings of the 2nd International Conference on Fifth Generation Computer Systems*, pages 85–99, 1984.
- [8] A. Colmerauer, H. Kanoui, and M. V. Caneghem. Prolog, theoretical principles and current trends. *Technology and Science of Informatics*, 2(4):255–292, 1983.
- [9] S. Haridi and S. Janson. Kernel Andorra Prolog and its computation model. In D. Warren and P. Szeredi, editors, *Logic Programming, Proceedings of the 7th International Conference*, pages 31–48, Cambridge, MA, June 1990. The MIT Press.
- [10] J. Jaffar and J.-L. Lassez. Constraint logic programming. In *Proceedings of the 14th ACM Symposium on Principles of Programming Languages*, pages 111–119, Munich, Germany, Jan. 1987.
- [11] M. Johnson. *Attribute-Value Logic and the Theory of Grammar*. CSLI Lecture Notes 16. Center for the Study of Language and Information, Stanford University, CA, 1988.
- [12] R. M. Kaplan and J. Bresnan. Lexical-Functional Grammar: A formal system for grammatical representation. In J. Bresnan, editor, *The Mental Representation of Grammatical Relations*, pages 173–381. The MIT Press, Cambridge, MA, 1982.

- [13] M. Kay. Functional grammar. In *Proceedings of the Fifth Annual Meeting of the Berkeley Linguistics Society*, Berkeley, CA, 1979. Berkeley Linguistics Society.
- [14] J.-L. Lassez, M. Maher, and K. Marriot. Unification revisited. In J. Minker, editor, *Foundations of Deductive Databases and Logic Programming*. Morgan Kaufmann, Los Altos, CA, 1988.
- [15] J. L. Lassez and K. McAloon. A constraint sequent calculus. In *Fifth Annual IEEE Symposium on Logic in Computer Science*, pages 52–61, June 1990.
- [16] M. J. Maher. Logic semantics for a class of committed-choice programs. In J.-L. Lassez, editor, *Logic Programming, Proceedings of the Fourth International Conference*, pages 858–876, Cambridge, MA, 1987. The MIT Press.
- [17] K. Mukai. Partially specified terms in logic programming for linguistic analysis. In *Proceedings of the 6th International Conference on Fifth Generation Computer Systems*, 1988.
- [18] K. Mukai. *Constraint Logic Programming and the Unification of Information*. PhD thesis, Tokyo Institute of Technology, Tokyo, Japan, 1991.
- [19] M. Nivat. Elements of a theory of tree codes. In M. Nivat, A. Podelski, editors, *Tree Automata (Advances and Open Problems)*, Amsterdam, NE, 1992. Elsevier Publishers.
- [20] W. C. Rounds and R. T. Kasper. A complete logical calculus for record structures representing linguistic information. In *Proceedings of the 1st IEEE Symposium on Logic in Computer Science*, pages 38–43, Boston, MA, 1986.
- [21] V. Saraswat and M. Rinard. Concurrent constraint programming. In *Proceedings of the 7th Annual ACM Symposium on Principles of Programming Languages*, pages 232–245, San Francisco, CA, January 1990.
- [22] G. Smolka. Feature constraint logics for unification grammars. *The Journal of Logic Programming*, 12:51–87, 1992.
- [23] G. Smolka and H. Aït-Kaci. Inheritance hierarchies: Semantics and unification. *Journal of Symbolic Computation*, 7:343–370, 1989.
- [24] G. Smolka and R. Treinen. Relative simplification for and independence of CFT. Draft, German Research Center for Artificial Intelligence (DFKI), Stuhlsatzenhausweg 3, 6600 Saarbrücken 11, Germany, 1992. To appear.